

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : A01N 1/02, C12Q 1/00 G01N 33/566, 33/543 C07D 471/02, 235/00, 473/00 C07D 235/30</p>	<p>A1</p>	<p>(11) International Publication Number: WO 91/07087</p> <p>(43) International Publication Date: 30 May 1991 (30.05.91)</p>
<p>(21) International Application Number: PCT/US90/06607</p> <p>(22) International Filing Date: 13 November 1990 (13.11.90)</p> <p>(30) Priority data: 435,316 13 November 1989 (13.11.89) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US Not furnished (CIP)</p> <p>(71) Applicant (for all designated States except US): AFFYMAX TECHNOLOGIES, N.V. [NL/NL]; De Ruyderkade 62, Curacao (AN).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): BARRETT, Ronald, W. [US/US]; 562 Carlisle Way, Sunnyvale, CA 94087 (US). PIRRUNG, Michael, C. [US/US]; 3421 Cottonwood, Durham, NC 27707 (US). STRYER, Lubert [US/US]; 843 Sonoma Terrace, Stanford, CA 94305 (US). HOLMES, Christopher, P. [US/US]; 521 Pine Street, Sunnyvale, CA 94086 (US). SUNDBERG, Steven, A. [US/US]; 659 Scott Street, #10, San Francisco, CA 94117 (US).</p>		<p>(74) Agent: SMITH, William, M.; Townsend and Townsend, One Market Plaza, 2000 Steuart Tower, San Francisco, CA 94105 (US).</p> <p>(81) Designated States: AT, AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CM (OAPI patent), DE, DE (European patent), DK, DK (European patent), ES, ES (European patent), FI, FR (European patent), GA (OAPI patent), GB, GB (European patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL, NL (European patent), NO, RO, SD, SE, SE (European patent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent), US.</p> <p>Published With international search report.</p>
<p>(54) Title: SPATIALLY-ADDRESSABLE IMMOBILIZATION OF ANTI-LIGANDS ON SURFACES</p> <p>(57) Abstract</p> <p>Methods and compositions are described for immobilizing anti-ligands, such as antibodies or antigens, hormones or hormone receptors, oligonucleotides, and polysaccharides on surfaces of solid substrates for various uses. The methods provide surfaces covered with caged binding members which comprise protecting groups capable of being removed upon application of a suitable energy source. Spatially addressed irradiation of predefined regions on the surface permits immobilization of anti-ligands at the activated regions on the surface. Cycles of irradiation on different regions of the surface and immobilization of different anti-ligands allows formation of an immobilized matrix of anti-ligands at defined sites on the surface. The immobilized matrix of anti-ligands permits simultaneous screenings of a liquid sample for ligands having high affinities for certain anti-ligands of the matrix. A preferred embodiment of the invention involves attaching photoactivatable biotin derivatives to a surface. Photolytic activation of the biotin derivatives forms biotin analogs having strong binding affinity for avidin. Biotinylated anti-ligands can be immobilized on activated regions of the surface previously treated with avidin.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	ML	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	SD	Sudan
CF	Central African Republic	KP	Democratic People's Republic of Korea	SE	Sweden
CG	Congo			SN	Senegal
CH	Switzerland	KR	Republic of Korea	SU	Soviet Union
CI	Côte d'Ivoire	LI	Liechtenstein	TD	Chad
CM	Cameroon	LK	Sri Lanka	TG	Togo
DE	Germany	LU	Luxembourg	US	United States of America
DK	Denmark	MC	Monaco		
ES	Spain	MG	Madagascar		

5

SPATIALLY-ADDRESSABLE IMMOBILIZATION
OF ANTI-LIGANDS ON SURFACES

BACKGROUND OF THE INVENTION

The present invention relates generally to methods
10 and compositions useful for immobilizing anti-ligands on
surfaces. The immobilized anti-ligands, which can be, for
example, hormones or hormone receptors, antibodies or antigens,
oligosaccharides, and oligonucleotides, may be used in a
variety of screening and assay methodologies for ligands in
15 liquid media.

Certain biological molecules are known to interact
and bind to other molecules in a very specific manner.
Essentially any molecules having a high binding specificity for
each other can be considered a ligand/anti-ligand pair, e.g., a
20 vitamin binding to a protein, a cell-surface receptor binding
to a hormone or drug, a glycoprotein serving to identify a
particular cell to its neighbors, an IgG-class antibody binding
to an antigenic determinant, an oligonucleotide sequence
binding to its complementary fragment of RNA or DNA, and the
25 like.

The specific binding properties of anti-ligands for
ligands have implications for many fields. For example, the
strong binding affinity of antibodies for specific determinants
on antigens is critical to the field of immunodiagnosics.
30 Additionally, pharmaceutical drug discovery, in many cases,
involves discovering novel drugs having desirable patterns of
specificity for naturally-occurring receptors or other
biologically important anti-ligands. Many other areas of
research exist in which the selective interaction of
35 anti-ligands for ligands is important and are readily apparent
to those skilled in the art.

The immobilization of anti-ligands onto surfaces is
an important step in performing repetitive assays and

screenings of ligands with solid phase systems. Previous methods of attaching anti-ligands to surfaces are limited by low reaction efficiencies or by a general inability to regionally and selectively attach a plurality of anti-ligands to the surface.

A large variety of methods are known for attaching biological molecules to solid supports. See generally, Affinity Techniques. Enzyme Purification: Part B. Methods in Enzymology, Vol. 34, ed. W.B. Jakoby, M. Wilchek, Acad. Press, NY (1974) and Immobilized Biochemicals and Affinity Chromatography, Advances in Experimental Medicine and Biology, vol. 42, ed. R. Dunlap, Plenum Press, NY (1974), which are incorporated herein by reference. For example, U.S. Patent No. 4,681,870 describes a method for introducing free amino or carboxyl groups onto a silica matrix. These groups may subsequently be covalently linked to, e.g., a protein or other anti-ligand, in the presence of a carbodiimide. Alternatively, a silica matrix may be activated by treatment with a cyanogen halide under alkaline conditions. The anti-ligand is covalently attached to the surface upon addition to the activated surface. Another example is presented in U.S. Patent No. 4,282,287, which describes a method for modifying a polymer surface through the successive application of multiple layers of biotin, avidin and extenders. Also, U.S. Patent No. 4,762,881 describes a method for attaching a polypeptide chain to a solid substrate by incorporating a light-sensitive unnatural amino acid group into the polypeptide chain and exposing the product to low-energy ultraviolet light.

Similarly, a variety of techniques have been developed for attaching oligonucleotides to surfaces. For example, U.S. Patent No. 4,542,102 describes a method employing a photochemically active reagent (e.g., a psoralen compound) and a coupling agent, which attaches the photoreagent to the substrate. Photoactivation of the photoreagent binds a nucleic acid sequence to the substrate to give a surface-bound probe for a complementary oligonucleotide of the sequence. However, this method has low quantum yields in protic solvents, lacks

spatial directability, and relies upon initial affinity between the photoreagent and nucleic acids prior to photoactivation.

U.S. Patent No. 4,562,157 describes a technique for attaching biochemical ligands to surfaces by attachment of a photochemically reactive arylazide. Irradiation of the azide creates a reactive nitrene which reacts irreversibly with macromolecules in solution resulting in the formation of a covalent bond. The high reactivity of the nitrene intermediate, however, results in both low coupling efficiencies and many potentially unwanted products due to nonspecific reactions.

Thus, there exists a need for improved methods for attaching a broad range of anti-ligands to predefined regions of a solid support surface. The methods should efficiently provide stable attachment of selected anti-ligands to the activated surface regions, yet attachment should be restricted to the activated regions. The present invention fulfills these and other needs.

SUMMARY OF THE INVENTION

Novel methods and compositions of matter are provided for immobilizing anti-ligands on predefined regions of a surface of a solid support. The methods involve attaching to the surface a caged binding member which has a relatively low affinity for other potentially binding species, such as anti-ligands and specific binding substances. The caged binding member is convertible, e.g., by irradiation, to a binding member ultimately capable of immobilizing a desired anti-ligand, preferably via a non-covalent interaction. Predefined regions of the surface are selectively irradiated to convert the caged binding members in the predefined regions to activated binding members. The desired anti-ligands subsequently can be immobilized on the activated regions of the surface.

Importantly, the spatial addressability afforded by the method of the present invention allows the formation of patterned surfaces having preselected reactivities. For example, by using lithographic techniques known in the

semiconductor industry, light can be directed to relatively small and precisely known locations on the surface. It is, therefore, possible to activate discrete, predetermined locations on the surface for attachment of anti-ligands. The resulting surface will have a variety of uses. For example, direct binding assays can be performed in which ligands can be simultaneously tested for affinity at different anti-ligands attached to the surface. Ligand binding is detected by a technique such as autoradiography when the ligand is radioactively labelled. Alternatively, fluorescence or other optical techniques can be used. By determining the locations and intensities of labels on the surface it is possible to simultaneously screen ligands for affinity to a plurality of anti-ligands.

A further understanding of the nature and advantages of the invention may be realized by reference to the remaining portions of the specification.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 presents chromatographic results showing that NVOC-biotin-ONP is converted to biotin-ONP upon illumination in solution.

FIG. 2 presents radioligand binding results showing NVOC-biotin-OMe has low affinity for avidin prior to illumination but high affinity after illumination in solution.

FIG. 3 presents gamma counting results showing that illumination of membrane-bound NVOC-biotin increases the binding of radioactive avidin to the membrane.

FIG. 4 presents fluorescence results showing the spatial immobilization of Fluorescein-Streptavidin on a biotinylated surface.

FIG. 5 presents fluorescence results showing the spatial immobilization of Fluorescein-Biotin on a surface modified with Streptavidin.

FIG. 6 presents fluorescence results showing the spatial immobilization of Fluorescein-Streptavidin on a surface having biotin bound by a polyether linker.

FIG. 7 presents fluorescence results showing the spatial immobilization of Fluorescein-Streptavidin on a biotinylated surface.

FIG. 8 presents fluorescence results showing the spatial immobilization of Bodipy-Streptavidin on a biotinylated surface.

FIG. 9 presents fluorescence results showing the effect of inserting linkers of different lengths on the binding of Fluorescein-Streptavidin.

FIG. 10a presents fluorescence results showing binding of Fluorescein-anti-Rabbit IgG to a slide having multiple anti-ligands.

FIG. 10b presents fluorescence results showing binding of Fluorescein-anti-Mouse IgG to a slide having multiple anti-ligands.

FIG. 10c presents fluorescence results showing binding of a mixture of Fluorescein-anti-Rabbit IgG and Fluorescein-anti-Mouse IgG to a slide having multiple anti-ligands.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS.

CONTENTS:

- I. Glossary
- II. Overview
- III. Substrate Preparation
- IV. Properties of Binding Members and Caging Groups
 - A. Caging Groups
 - B. Irradiation of Caged Compounds
- V. Attachment of Anti-Ligands
- VI. Screenings and Assays
- VII. Examples

I. Glossary

The following terms have the following meanings and abbreviations as used herein:

1. Surface (S): A surface is any generally two-dimensional structure on a solid substrate. A surface may have steps, ridges, kinks, terraces and the like without ceasing to be a surface.

2. Predefined Region (S_1): A predefined region is a localized area on a surface which is or is intended to be

activated. The predefined region may have any convenient shape, e.g., circular, rectangular, elliptical, etc.

5 3. Crosslinking Group (X): A crosslinking group is a bifunctional chemical entity that serves to connect a binding member to a surface. Usually, crosslinking groups will be heterobifunctional, i.e., they will have different chemical reactivities on either end of the linking group.

10 4. Binding Member (B): A binding member is any substance having a sufficiently high affinity for another substance. A binding member will have a sufficiently high affinity for another substance for practice of this invention when it effectively binds the substance without irreversibly separating from it throughout the handling and performance steps of the invention. A binding member is usually, but not
15 always, connected to a surface via a crosslinking group.

5. Caged Binding Member (B*): A caged binding member is a binding member that is provided with a removable (labilizable) chemical protecting group. Such protecting groups are characterized by their abilities to deter
20 effectively binding between the binding member to which they are attached and other substances otherwise having affinity for the binding member. Also, the protecting groups are readily labilizable, i.e., they can be detached from the binding member to which they are attached upon exposure to a suitable source
25 of energy.

6. Specific Binding Substance (SBS): A specific binding substance is a compound having a sufficiently high affinity and selectivity for binding to a binding member to permit practice of the present invention. A specific binding
30 substance may be larger or smaller than the binding member to which it specifically binds. The specific binding substance serves as a bridge for attaching an anti-ligand to binding members on the surface.

7. Anti-ligand (AL_i): An anti-ligand is a molecule
35 that has a known or unknown affinity for a given ligand and can be immobilized on a predefined region of the surface. Anti-ligands may be naturally-occurring or manmade molecules. Also, they can be employed in their unaltered state or as

aggregates with other species. Anti-ligands may be reversibly attached, covalently or noncovalently, to a binding member, either directly or via a specific binding substance. By "reversibly attached" is meant that the binding of the anti-ligand (or specific binding member or ligand) is reversible and has, therefore, a substantially non-zero reverse, or unbinding, rate. Such reversible attachments can arise from noncovalent interactions, such as electrostatic forces, van der Waals forces, hydrophobic (i.e., entropic) forces, and the like. Furthermore, reversible attachments also may arise from certain, but not all covalent bonding reactions. Examples include, but are not limited to, attachment by the formation of hemiacetals, hemiketals, imines, acetals, ketals, and the like (See, Morrison et al., "Organic Chemistry", 2nd ed., ch. 19 (1966), which is incorporated herein by reference). Examples of anti-ligands which can be employed by this invention include, but are not restricted to, cell membrane receptors, monoclonal antibodies and antisera reactive with specific antigenic determinants (such as on viruses, cells or other materials), hormones, drugs, oligonucleotides, peptides, enzymes, substrates, cofactors, lectins, sugars, oligosaccharides, cells, cellular membranes, and organelles.

8. Ligand (L): A ligand is a solvated molecule that is recognized by a particular anti-ligand. Examples of ligands that can be investigated by this invention include, but are not restricted to agonists and antagonists for cell membrane receptors, toxins and venoms, viral epitopes, hormones (e.g., opiates, steroids etc.), hormone receptors, peptides, enzymes, enzyme substrates, cofactors, drugs, lectins, sugars, oligonucleotides, oligosaccharides, proteins, and monoclonal antibodies.

II. Overview

The present invention provides methods for forming predefined regions on a surface of a solid support, wherein the predefined regions are capable of immobilizing anti-ligands. The methods make use of caged binding members attached to the surface to enable selective activation of the predefined

regions. The caged binding members are converted to binding members ultimately capable of binding anti-ligands upon selective activation of the predefined regions. The activated binding members are then used to immobilize anti-ligands on the predefined region of the surface. The above procedure can be repeated at the same or different sites on the surface so as to provide a surface prepared with a plurality of regions on the surface containing the same or different anti-ligands. When the anti-ligands have a particular affinity for one or more ligands, screenings and assays for the ligands can be conducted in the regions of the surface containing the anti-ligands.

The present methods are distinguished by the employment of novel caged binding members attached to the substrate. Caged (unactivated) members have a relatively low binding affinity for anti-ligands or specific binding substances when compared with the corresponding affinities of activated binding members. Thus, the binding members are protected until a suitable source of energy is applied to the regions of the surface desired to be activated. Upon application of a suitable energy source, the caging groups labilize, thereby presenting the activated binding member. A typical energy source will be light.

Once the binding members on the surface are activated they may be attached to an anti-ligand. The anti-ligand chosen may be a monoclonal antibody, a nucleic acid sequence, a drug receptor, etc. The anti-ligand will usually, though not always, be prepared so as to permit attaching it, directly or indirectly, to a binding member. For example, a specific binding substance having a strong binding affinity for the binding member and a strong binding affinity for the anti-ligand may be used as a bridge. Alternatively, a covalently-linked conjugate of the specific binding substance and anti-ligand may be used. The method uses an anti-ligand prepared such that the anti-ligand retains its activity toward a particular ligand.


Preferably, the caged binding member attached to the solid substrate will be a photoactivatable biotin analog, i.e., a biotin molecule that has been chemically modified with

photoactivatable protecting groups so that it has a significantly reduced binding affinity for avidin or avidin analogs compared to that of natural biotin. In a preferred embodiment, the protecting groups localized in a predefined region of the surface will be removed upon application of a suitable source of radiation to give binding members, that are biotin or a functionally analogous compound having substantially the same binding affinity for avidin or avidin analogs as does biotin.

In another preferred embodiment, avidin or an avidin analog will be incubated with activated binding members on the surface until the avidin binds strongly to the binding members. The avidin so immobilized on predefined regions of the surface, can then be incubated with a desired anti-ligand or conjugate of a desired anti-ligand. The multiple biotin binding sites on avidin allow simultaneous binding of biotin attached to the surface and biotin attached to the anti-ligand. The anti-ligand will preferably be biotinylated, e.g., a biotinylated antibody, when avidin is first immobilized on the predefined regions of the surface. Alternatively, a preferred embodiment will present an avidin/biotinylated anti-ligand complex, which has been previously prepared, to activated binding members on the surface.


The following equations depict the best modes of practicing the invention:

The attachment of binding members (B) to a surface (S) of a solid substrate is illustrated by the following reactions:




where "*" represents a protecting (caging) group. B* is a caged binding member. The protecting groups can either be attached to the binding members once they have been attached to the surface, or more preferably, they will be attached to binding members prior to attaching the binding members to the surface.

Also, surface attachment of binding members can be effected through the use of crosslinking groups (X). This is represented by the following reactions:



The crosslinking groups will usually, though not always, be heterobifunctional chemical species having a first reactivity which permits the crosslinking group to bind readily to the surface and a second reactivity which permits the crosslinking group to bind readily with binding members.


Predefined regions (S_i) on the surface can be activated for ultimate immobilization of anti-ligands in the predefined regions by selectively irradiating predefined regions to convert photoactivatable binding members in the predefined region to binding members. This process is illustrated by the following reactions:



The free protecting group, "*", may or may not undergo decomposition reactions. It will usually be washed from the surface, depending upon whether it interferes with subsequent reactions.


Immobilization of anti-ligands (AL_i) on predefined regions of the surface can be effected by binding the anti-ligands directly to binding members or through a bridging specific binding substance (SBS). The specific binding substance may be introduced to binding members alone or as a previously prepared conjugate of the anti-ligand. Multiple anti-ligands may be immobilized on the surface when the specific binding substance contains multiple binding sites. Also, it should be noted that an advantage of using a specific binding substance is that an immobilization technique generic for many anti-ligands may be employed. Immobilization of

anti-ligands on predefined regions of the surface is illustrated by the following reactions:




where the horizontal lines B-SBS, B-AL_i, or SBS-AL_i represent bonding between two molecules, preferably a non-covalent bond.


An example of immobilizing a different anti-ligand (AL_j) on a different predefined region (S_j) of the surface is shown by the equation:



Repetition of the above steps on different regions of the surface can produce a matrix of anti-ligands immobilized on the surface. Such a matrix can have any desired pattern of anti-ligands. An example of such a matrix is given below:




An immobilized anti-ligand on a surface will have a specific binding affinity for a particular ligand (L). An example of a direct assay on a predefined region of the surface for the presence of a labeled ligand (L') in a liquid medium is illustrated by the following reaction:



The resulting surface can be washed free of unbound ligand and analyzed for the presence of label. The labels will provide markers localized at the predefined regions on the surface corresponding to the presence of anti-ligands for the ligand at those predefined regions.

Some examples of competitive assays, in which a target ligand (L) "competes" with another ligand (L') for a site on the surface, are illustrated by the following reactions:

5



10 The presence of target ligand can be determined by analyzing appropriately for the loss or buildup of label on the predefined regions of the surface.

III. Substrate Preparation

Essentially, any conceivable solid substrate may be employed in the invention. The substrate may be biological, nonbiological, organic, inorganic, or a combination of any of these, existing as particles, strands, precipitates, gels, sheets, tubing, spheres, containers, capillaries, pads, slices, films, plates, slides, etc. The substrate may have any convenient shape, such as a disc, square, sphere, circle, etc. The substrate and its surface preferably form a rigid support on which to carry out the reactions described herein. The substrate and its surface should also be chosen to provide appropriate light-absorbing characteristics. For instance, the substrate may be a polymerized Langmuir Blodgett film, functionalized glass, Si, Ge, GaAs, GaP, SiO₂, SiN₄, modified silicon, or any one of a wide variety of polymers such as (poly)tetrafluoroethylene, (poly)vinylidenedifluoride, or combinations thereof. Other substrate materials will be readily apparent to those of skill in the art upon review of this disclosure. In a preferred embodiment the substrate is flat glass or single-crystal silicon with surface features of less than 10 Å.

Surfaces on the solid substrate will usually, though not always, be composed of the same material as the substrate. Thus, the surface may be composed of any of a wide variety of materials, for example, polymers, plastics, resins, polysaccharides, silica or silica-based materials, carbon, metals, inorganic glasses, membranes, etc., provided only that

caged binding members can be attached firmly to the surface of the substrate. Preferably, the surface will contain reactive groups, which could be carboxyl, amino, hydroxyl, or the like. Most preferably, the surface will be optically transparent and
5 will have surface Si-OH functionalities, such as are found on silica surfaces.

The surface of the substrate is preferably provided with a layer of crosslinking groups, although it will be understood that the crosslinking groups are not required
10 elements of the invention. The crosslinking groups are preferably of sufficient length to permit binding members on the surface to interact freely with compounds in solution. Crosslinking groups may be selected from any suitable class of compounds, for example, aryl acetylenes, ethylene glycol
15 oligomers containing 2-10 monomer units, diamines, diacids, amino acids, or combinations thereof. Other crosslinking groups may be used in light of this disclosure.

Crosslinking groups may be attached to the surface by a variety of methods which are readily apparent to one having
20 skill in the art. For example, crosslinking groups may be attached to the surface by siloxane bonds formed via reactions of crosslinking groups bearing trichlorosilyl or trisalkoxy groups with hydroxyl groups on the surface of the substrate. Preferably, the crosslinking group used with a glass surface is
25 N-BOC-aminopropyltriethoxy silane. The crosslinking groups may optionally be attached in an ordered array, i.e., as parts of the head groups in a polymerized Langmuir Blodgett film. Clearly, the type of crosslinking group selected, and the method selected for attaching it to the surface, will depend
30 primarily on the crosslinking group having suitable reactivity with the binding member desired to be attached to the surface.

Additional length may be added to the crosslinking groups by the addition of single or multiple linking groups. Such linking groups are preferably heterobifunctional, having
35 one end adapted to react with the crosslinking groups and the other end adapted to react with the binding member or an another linking group. The linking groups may be attached by a variety of methods which are readily apparent to one skilled in

the art. For instance, esterification or amidation reactions of an activated ester of the linking group with a reactive hydroxyl or amine on the free end of the crosslinking group. A preferred linking group is N-BOC-6-aminocaproic acid (i.e., N-BOC-6-aminohexanoic acid) attached by the BOP-activated ester. After deprotection to liberate the free amine terminus, another N-BOC-aminocaproic linker can be added. Attachment of crosslinking and linking groups to caged binding members are discussed more fully below.

Many methods are available for immobilizing the binding members of the present invention on surfaces. The binding members may be linked to the surface in their active forms, and later provided with protecting (caging) groups. More preferably, binding members will be provided in their protected forms. The method chosen for linking binding members to the surface will depend upon the chemical properties of the binding member selected for attachment to the surface. A preferred method for immobilizing the binding members of the present invention involves chemical derivatization or activation of the caged binding member prior to attachment to the surface or linker. This derivative or activated species is then reacted with functionalities on the substrate to give the desired linkage. For example, one method for attaching a binding member to a surface employs a heterobifunctional crosslinking reagent, such as diepoxide, which both activates the surface and provides a group that reacts with an activated binding member. Alternatively, the surface can be activated with cyanogen bromide. Reaction with a binding member containing a terminal amino group permits attachment of the binding member to the surface. (U.S. Patent No. 4,542,102). In the presence of a carbodiimide or other activating agent, for example, the amine group can be coupled to the carboxyl terminus of a binding member desired to be immobilized on the surface.

A preferred embodiment of the present invention involves attaching "caged" derivatives of biotin or biotin analogs to a glass surface. Caged biotin may be attached to the surface through strong noncovalent interactions, e.g., by

crosslinking via a suitable linker to another biotin molecule and reacting with a surface to which avidin has been attached, or alternatively, and preferably, by covalent attachment to the surface. The latter may be accomplished by derivatizing caged-
5 biotin and biotin analogues at their carboxylic acid terminus. Many biotin derivatives have been described previously. For example, the surface can be provided with biotin anti-ligands, e.g., antibiotin antibodies, which specifically bind the carboxyl arm of biotin without interfering with the
10 avidin-binding ureido ring of biotin.

Still another method for immobilizing the caged binding members of the present invention involves chemical derivatization or activation of the binding member prior to attachment to the surface or linker. For example, when the
15 surface is a polymer containing primary amines and biotin is selected as the binding member, the N-hydroxysuccinimide ester derivative of biotin can react with the surface to give a biotin-surface complex (U.S. Patent No. 4,282,287).

Alternatively, and preferably, photoactivatable
20 biotin and biotin analog derivatives will be covalently attached to the surface. To effect this transformation, the biotin and biotin analogs may be derivatized at their carboxylic acid terminus. Many biotin derivatives have been described previously involving derivatization at the free
25 carboxyl end of biotin. See, e.g., Bayer et al., Methods of Biochemical Analysis, vol. 26 (D. Glick, ed.), 1-45 (1980), which is incorporated herein by reference. For example, photoactivatable biotin derivatives may be reacted, in the presence of an activating reagent, such as a carbodiimine or
30 BOP, with the amine groups of crosslinking groups previously immobilized on the surface to give the biotin attached to the surface via an amide linkage. The active ureido ring of biotin, either free or protected, is located far enough away from the site of attachment that, when unprotected, binding
35 with avidin is not significantly diminished.

It should be appreciated that the above discussion of exemplary surface attachment reactions is only illustrative of the general method for attaching caged binding members to a

surface and should not be regarded in any way as limiting the applicability of the method to biotin, biotin analogs, or specific crosslinking groups. Other types of binding members also amenable to the above attachment techniques include enzymes, antibodies, oligonucleotides and the like and are readily apparent to one skilled in the art.

IV. Properties of Binding Members and Caging Groups

The present method permits use of a wide variety of caged binding members to effect the immobilization of anti-ligands on the surface. The method is generally applicable to such classes of compounds as enzymes, substrates, cofactors, immunoglobulins, antibodies, haptens, antigens, oligonucleotides, oligosaccharides, lectins, proteins, glycoproteins, etc., being the binding member provided that the selected derivative of such species is activatable upon exposure to a suitable energy source. Moreover, the binding member can possess a multiplicity of binding sites for an anti-ligand or specific binding substance.

The binding member selected will have a high binding affinity either for an anti-ligand or a specific binding substance. Preferably, a specific binding substance will provide a link between the binding member and the anti-ligand. Usually, the interactions between a binding member and a specific binding substance and an anti-ligand or anti-ligand conjugate will be noncovalent in nature. When a specific binding substance provides a link between the binding member and the anti-ligand, the specific binding substance will be connected to the anti-ligand either covalently or through noncovalent interactions.

The binding member on a surface must have a strong affinity for an anti-ligand or specific binding substance to prevent migration or loss of the anti-ligand during wash steps. The affinity between the binding member and an anti-ligand or a specific binding substance is best represented by the off-rate of anti-ligand or specific binding substance from a binding member. However, off-rates often are not conveniently known or determined. Therefore, binding affinity may also be

represented by the affinity constant (K_a) for equilibrium concentrations of associated and dissociated configurations; i.e., $K_a = [B-SBS]/[B][SBS]$, where $[B]$, $[SBS]$ and $[B-SBS]$ are the concentrations of the binding member (B), the concentration of specific binding substance (SBS), and the concentration of associated complex (B-SBS), respectively, at equilibrium. An analogous definition of K_a applies when SBS is replaced with an anti-ligand (AL) or a specific binding substance - anti-ligand conjugate (SBS-AL), etc. The affinity constants of some sample classes of compounds suitable for use in the present invention are presented in Table 1.

Table 1

Affinities of Sample Binding Members and Specific Binding Substances (SBS).

<u>Binding Member</u>	<u>SBS</u>	<u>Affinity (K_a M⁻¹)</u>
Membrane sites	Lectins	10 ⁶⁻⁷
Haptens	Antibodies	10 ⁵⁻¹¹
Antigenic determinants	Antibodies	10 ⁵⁻¹¹
Biotin	Avidin	10 ¹⁵
Iminobiotin	Avidin	10 ¹¹
2-thiobiotin	Avidin	10 ¹³
Dethiobiotin	Avidin	10 ¹³
1'-N-methoxy-carbonylbiotin methyl ester	Avidin	10 ⁷
3'-N-methoxy-carbonylbiotin methyl ester	Avidin	10 ⁹

*References: U.S. Patent No. 4,282,287; Green, "Avidin" in Advances in Protein Chemistry, Academic Press, vol.29, 105 (1975).

Preferably, the affinity constant between the activated binding member and another species, i.e., a specific binding species, an anti-ligand, or anti-ligand conjugate, will be greater than about 10⁷ M⁻¹. More preferably, the K_a will be greater than about 10¹¹ M⁻¹, and most preferably, the K_a will be about 10¹⁵ M⁻¹ or greater. Likewise, when a specific binding substance is used, the affinity constant between the specific binding substance and an anti-ligand or anti-ligand conjugate will have substantially the same ranges as given above.

An activated (uncaged) binding member is considered to have a relatively strong (high) binding affinity for another species, i.e., a specific binding substance, an anti-ligand, or a conjugate of an anti-ligand, when the K_a between the binding member and the other species is at least about three orders of magnitude greater than the corresponding K_a between the caged binding member and the other species. Similarly, a caged binding member is considered to have a relatively low binding affinity for another species, i.e., specific binding substance, anti-ligand or anti-ligand conjugate, when the K_a between the caged binding member and the other species is about three orders of magnitude less than the corresponding K_a for the activated binding member. Preferably, the affinity constant for the caged binding member will be at least five orders of magnitude lower than the corresponding activated binding member's affinity constant. Most preferably, the binding constant for the caged binding member will be even lower, e.g., seven orders of magnitude lower, than the corresponding activated binding member's affinity constant. However, the suitability of a given caged binding member/ binding member pair for practice of the invention is determined ultimately by whether the selected pair permits proper operation of the invention.

A preferred embodiment of the present invention employs biotin and biotin analogs as the binding members. Typical biotin analogs include dethiobiotin, iminobiotin, 2-thiobiotin, azabiotin, biocytin, and biotin sulfone, and other compounds readily apparent to one skilled in the art. Exemplary biotin analogs include, but are not limited by, those presented in Table 2. Other biotin analogs are presented in N. Green, "Avidin", in Advances in Protein Chemistry, Vol. 29, Acad. Press, p. 85-133 (1975), which is incorporated by reference herein. Biotin analogs include compounds and structures in which biotin is bound to another species, such as a surface, as long as the analog has a binding affinity for avidin that is similar to that of biotin. The biotin or biotin analogs may be subsequently reacted with avidin compounds, streptavidin, and analogues thereof.

Typical examples of avidin and avidin analogs include, but are not limited to, the avidin found in eggs and streptavidin. Streptavidin is a typical example of an avidin analog and is a bacterial biotin-binding protein which has
5 physical characteristics similar to those of egg avidin, despite considerable differences in composition.

Table 2

Biotin and Biotin Analogs	Name
	Biotin
	Iminobiotin
	Dethiobiotin
	2'-thiobiotin
	Azabiotin
	Bisnorazabiotin

55 Reference: N. Green, "Avidin," in Advances in Protein Chemistry, Vol. 29, Academic Press, p. 85-133 (1975).

Nonbiotin binding members and their corresponding specific binding substances may be employed in the present invention. By way of example and not limitation, some alternative embodiments of the invention follow:

5

1. Caged Cyclic AMP/ Anti-cAMP Antibodies

High-affinity polyclonal antibodies to cAMP are produced by immunizing with 2'O-monosuccinyl adenosine 3',5' cyclic monophosphate conjugated to a protein such as bovine serum albumin or thyroglobulin. Purified polyclonal antibodies are prepared by affinity chromatography using agarose gel to which 2'O-monosuccinyl adenosine 3',5' cyclic monophosphate has been conjugated. The K_a of polyclonal antibodies to cAMP is in the range of 10^{10} to 10^{12} M^{-1} .

15

A photoactivatable analog of cAMP has been previously described (Nerbone et al., Nature (1984) 310:74). It is unlikely that polyclonal antibodies against cAMP have high affinity for the photoactivatable analog of cAMP. If the polyclonal antibodies should cross-react with the photoactivatable analog of cAMP, monoclonal antibodies can be produced which discriminate between cAMP and the photoactivatable cAMP analog.

20

The 2'O-monosuccinyl derivative of the photo-activatable cAMP analog is attached to a surface through the free carboxyl of the succinyl group as described above. Specific regions of the surface are illuminated resulting in the removal of the protecting group from the cAMP. Anti-ligands which have been conjugated to anti-cAMP antibodies are reacted with the surface. The anti-ligands are immobilized only at the predefined regions of the surface that were illuminated.

30

2. Caged Tetrahydrofolate/ Folate Binding Proteins

N_5 -(Nitroveratryloxycarbonyl)tetrahydrofolate is activated at its glutamyl gamma-carboxylate with a carbodiimide reagent and coupled to an amino-derivatized surface. Desired predefined regions on the surface are irradiated with light suitable for deprotection of the NVOC group. In the

35

illuminated regions, the NVOC group is removed to produce tetrahydrofolate bound to the surface. High-affinity folate binding proteins derived from human erythrocyte membranes (Antony et al., J. Clin. Invest. (1987) 80:711-723;

- 5 $K_a = 3 \times 10^{11} \text{ M}^{-1}$ for tetrahydrofolate), crosslinked to a desired anti-ligand, are then immobilized on the selected regions of the surface.

3. Caged Mannose/ Concanavalin A

10 8'-(Trichlorosilyl)octyl

- 6-(nitroveratryloxy)- α -D-mannoside is covalently attached to a silica or glass surface by methods well-known to those skilled in the art. Predefined regions of the surface are irradiated with light suitable for deprotection of the nitroveratryloxy group. In the irradiated regions, the protecting group is removed to produce octyl- α -D-mannoside bound to the surface. In unexposed areas, the 6-(nitroveratryloxy) group protects the mannoside from binding to the lectin. Concanavalin A, crosslinked to a desired anti-ligand, is added to the surface and binds to those mannose units on the surface that have been deprotected. The anti-ligand is thereby immobilized on the desired regions of the surface.

- The above modes for practicing the invention are examples only. Those skilled in the art will recognize that any pair of (i) binding member and specific binding substance, (ii) binding member and anti-ligand or anti-ligand conjugate, or (iii) specific binding substance and anti-ligand or anti-ligand conjugate may be used. The only restrictions on the choice of binding member, specific binding substance and anti-ligand or anti-ligand conjugate are that: (1) the binding member has a high affinity for the specific binding substance, anti-ligand or anti-ligand conjugate selected, (2) the binding member can be "caged" with a removable protecting group, and (3) the caged binding member has a relatively low affinity for specific binding substances, anti-ligands, or anti-ligand conjugates, or any other species which interfere with practice of the invention.

A. Caging Groups

Many different protecting (caging) groups can be employed for modifying binding members to give the caged binding members of the present invention. The protecting groups should be sterically large enough to reduce the affinity of the binding member for anti-ligands or specific binding substances to permit operability of the invention, although protecting groups utilizing other types of interactions, such as electronic (i.e., Van der Waals), hydrophobic, etc., could be used. The selection of suitable caging groups will depend upon the size and chemical nature of the binding member chosen and will be readily apparent to one skilled in the art.

In a preferred embodiment, the caging groups will be photoactivatable. The properties and uses of photoreactive caged compounds have been reviewed. See, J. McCray, et al., Annu. Rev. Biophys., Biophys. Chem., 18: 239-70 (1989), which is incorporated herein by reference. Preferably, the photosensitive cages will be activatable by low energy ultraviolet or visible light. In some embodiments, however, activation may be performed by the methods discussed later, including localized heating, electron beam techniques, laser pumping, and oxidation or reduction with microelectrodes. Alternatively, the reactive group may be activatable by electron beam lithography, X-ray lithography, or any other radiation. Suitable reactive groups for electron beam lithography include sulfonyl compounds. Other methods may be used including, for example, exposure to an electric current source, preferably using microelectrodes directed to the predefined regions of the surface which are desired for activation. Other reactive groups and methods of activation may be used in light of this disclosure.


A further preferred embodiment of the present invention employs photoactivatable N-derivatives of biotin and biotin analogs to reduce the natural affinity of biotin for other compounds, such as avidin used as a specific binding substance, until the groups attached to the N-positions are photoremoved. A few references describe N-derivatization of biotin and biotin analogs. See, Kohn et al., J. Org. Chem.

(1977) 42:941-948, and Knappe et al., Biochemische Zeitschrift, 335, 168-176 (1961). However, none of these references provide photoactivatable biotin derivatives. The use of a photosensitive biotin derivative, photobiotin, has been previously described for labelling proteins and nucleic acids. Lacey, E. Anal. Biochem., 163: 151-8 (1987); Forster, A.C. Nucleic Acids Res., 13: 745-61 (1985). However, photobiotin is a derivative of the carboxylate terminus of biotin, which is located away from the recognition site and, hence, does not significantly reduce binding to avidin or streptavidin.


Many, although not all, of the photosensitive protecting groups will be aromatic compounds. Suitable photoremovable protecting groups are described in, for example, McCray, et al., Patchornik, J. Am. Chem. Soc. (1970) 92:6333 and Amit, et al., J. Org. Chem. (1974) 39:192, which are incorporated herein by reference. See, also, Calbiochem Catalog, (San Diego, CA, 1989), p. 244-247. More preferably, the photosensitive group will be a nitro benzylic compound, such as o-nitrobenzyl or benzylsulfonyl groups. In a preferred embodiment, 6-nitroveratryloxycarbonyl (NVOC) and its derivatives, such as and 6-nitropiperonyloxycarbonyl (NPOC), α,α -dimethyl-dimethoxybenzyloxycarbonyl (DDZ) or 1-pyrenylmethyl may be employed.

When the selected binding member is biotin or a biotin analog, photosensitive protecting groups may be provided at the N-1', N-3' or to an oxygen, imino, or sulfur group at the 2'-C position of the ureido ring. When the protecting group is attached to a 2'-C-O- position, the group will preferably be an o-nitro benzylic group having a hydrogen atom at the alpha benzylic position. In such case, the biotin or biotin residue is an imidazolidine group. When the protecting group is attached to a 1'-N or 3'-N atom, the protecting group will preferably be an o-nitro benzylic group having a hydrogen atom bound to the alpha carbon atom and optionally an oxycarbonyl group linking the alpha carbon atom through the oxygen atom. In the latter case, the derivatized nitrogen atom of the imidazolidone group will be bound to the carbon atom of the oxycarbonyl group.


A preferred embodiment of the invention has the following formula:



- 5 where X and Z are hydrogen, or oxycarbonyls of lower alkyl, aryl, or benzyl groups, provided that X and Z are not both hydrogen; R is hydrogen, lower alkyl, aryl, carboxylate, alkyl formate, aryl formate, formamide, N-alkylformamide, N-succinimidyl, hydroxyl, alkoxyl, thiol, thioether, disulfide, 15 hydrazide or an amine group; U is O, S, or NH; Y is sulfur, oxygen, methylene, carbonyl, or a sulfinyl, or sulfonyl group, or Y represents two hydrogen atoms attached to the respective carbons; and n = 0-7. Also, inorganic and organic acid addition salts of the above compounds can be employed.
- 20 Furthermore, R can represent a surface or a surface provided with a suitable crosslinking group. In another preferred embodiment; R is hydrogen, lower alkyl, aryl, carboxylate, alkyl formate, aryl formate, formamide, N-alkylformamide, N-succinimidyl, hydroxyl, alkoxyl, thiol, thioether, disulfide, 25 hydrazide or an amine group connected to a linking group of a suitable length, such as monomer, dimer, trimer, or oligomer of 6-aminocaproic acid, an oligomer of ethylene glycol having up to 10 units dioxadodecane-propyl, or other suitable linkers. A more preferred embodiment is when R is methyl formate or 30 p-nitrophenyl formate. A further preferred embodiment is when U is O, Y is S, and n=4. A preferred embodiment is when Y represents two hydrogen atoms attached to respective carbons, which eliminates the lower ring leaving a methyl group and a hexanoic acid group:



A further preferred embodiment is when X or Z is a nitro aromatic compound containing a benzylic hydrogen ortho to the nitro group. A still further preferred embodiment is when X or Z has the formula:




where R_1 and R_2 are hydrogen, lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxyl, thiol, thioether, amino, nitro, carboxyl, formate, sulfonate, formamido or phosphido groups.

A further preferred embodiment is when X or Z is a nitroveratryloxycarbonyl group.


Most preferred embodiments are when X is 6-nitroveratryloxycarbonyl, Z is hydrogen, and R is methyl formate or p-nitrophenyl formate.

Another preferred embodiment is when X or Z is a ring-disubstituted benzyloxycarbonyl group having the formula:



where R_1 and R_2 are hydrogen, lower alkyl, aryl, benzyl, pyrenyl, halogen, hydroxyl, alkoxyl, thiol, thioether, amino, nitro, carboxyl, formate, formamido or phosphido groups, and R_3 and R_4 are hydrogen, lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxyl, thiol, thioether, amino, nitro, carboxyl, formate, formamido or phosphido groups. More preferably R_1 and R_2 are methoxy groups. More preferably R_3 and R_4 are methyl groups. A most preferred embodiment is when R_1 and R_2 are methoxy groups and R_3 and R_4 are methyl groups.


A further preferred embodiment is when X has the formula:



where R_1 and R_2 are hydrogen, lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxyl, thiol, thioether, amino, nitro, carboxyl, formate, formamido or phosphido groups. A still further preferred embodiment is when X is a nitroveratryl group.


Most preferred embodiments are when X is 6-nitroveratryl, U or W is hydrogen, and R is methyl formate or p-nitrophenyl formate.

A further preferred embodiment is when X is a ring-disubstituted benzyl group. A more preferred embodiment is when the ring-disubstituted benzyl group has the formula:



where R_1 , R_2 , R_3 , and R_4 are hydrogen, lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxyl, thiol, thioether, amino, nitro, carboxyl, formate, formamido or phosphido groups. Most preferred embodiments are when R_1 and R_2 are methoxy and R_3 and R_4 are methyl.

Another preferred embodiment is when the composition has the following formula:



where X is hydrogen, lower alkyl, aryl, or benzyl; U and W are hydrogen, lower alkyl, aryl, or benzyl groups, provided that only one of U and W is present; R is hydrogen, lower alkyl, aryl, carboxylate, alkyl formate, aryl formate, formamide, N-alkyl formamide, N-succinimidyl, hydroxyl, alkoxyl, thiol, thioether, disulfide, hydrazide, or amine groups; Y is sulfur, oxygen, methylene, carbonyl, or a sulfinyl, or sulfonyl group, or Y represents two hydrogen atoms attached to the respective carbons; and $n = 0-7$. Also, inorganic and organic acid addition salts of the above compounds are suitable. A more preferred embodiment is when Y is sulfur and $n = 4$.

Clearly, many photosensitive protecting groups are suitable for use in the present inventive methods. Some examples of acceptable photosensitive protecting groups are presented in Table 3. Also, some example protecting groups and their corresponding wavelengths for deprotection are provided in Table 4.

Table 3







Example	Protecting Groups	Name
5		6-nitroveratryloxy-carbonyl (NVOC)
10		
15		dimethyldimethoxy-benzyloxycarbonyl (DDZ)
20		
25		nitrobenzyloxycarbonyl (NBOC)
30		
35		5-bromo-7-nitroindolinyl (BNI)
40		
45		o-hydroxy-α-methyl-cinnamoyl (HMC)
50		
55		2-oxymethylene anthraquinone (OMA)

Table 4:

<u>Example protecting groups and their deprotection wavelengths</u>	
<u>Group</u>	<u>Deprotection Wavelength</u>
5 Nitroveratryloxycarbonyl	UV (300-350 nm)
Nitrobenzyloxycarbonyl	UV (300-350 nm)
Dimethyldimethoxybenzyloxycarbonyl	UV (280-300 nm)
5-Bromo-7-nitroindoliny	UV (420 nm)
o-Hydroxy- α -methyl cinnamoyl	UV (300-350 nm)
10 2-Oxymethylene anthraquinone	UV (350 nm)

B. Irradiation

Once the surface is covered with a plurality of caged
 15 binding members, selected regions of the surface may be
 irradiated to provide activated binding members. Predefined
 regions of the surface may be selectively activated by electron
 beam lithography, ion beam lithography, X-ray lithography, or
 any other radiation method. In a preferred embodiment, the
 20 radiation is UV, near IR, or visible light. The light source
 may be coherent or noncoherent. The protective group may
 alternatively be an electrochemically-sensitive group which may
 be removed in the presence of an electric current.

In some embodiments, the exposed area is less than
 25 about 1 cm² or less than about 1 mm². In preferred embodiments
 the exposed area is less than about 10,000 μ m² or, more
 preferably, less than about 100 μ m². Spaces between activated
 regions are not critical and will generally be greater than
 about 1 μ m.

30 When photoactivatable binding members are used, they
 are preferably exposed to light through a suitable mask using
 photolithographic techniques well known in the semiconductor
 industry and described in, for example, Sze, VLSI Technology,
 McGraw-Hill (1983), which is incorporated herein by reference.
 35 In one embodiment, the mask is a transparent support material
 coated with a layer of opaque material. Portions of the opaque
 material are removed, leaving opaque material in the precise
 pattern desired on the substrate surface. The mask is brought
 into close proximity with or directly into contact with the
 40 surface. Openings in the mask correspond to locations on the
 surface where it is desired to photoremove protecting groups
 from the binding members. Alignment may be performed using

conventional alignment techniques in which alignment marks are used to accurately overlay successive masks with previous patterning steps. Other alignment techniques may be used, for example, interferometric techniques such as the one described in Flanders, et al., "A New Interferometric Alignment Technique," App. Phys. Lett. (1977) 31:426-428, which is incorporated herein by reference.

To enhance contrast of light applied to the substrate it may be desirable to provide contrast enhancement materials between the mask and the substrate. This contrast enhancement layer may comprise a molecule which is decomposed by light such as quinone diazide.

The light may be from a conventional incandescent source, an arc lamp, a laser, or the like. If noncoherent sources of light are used it may be desirable to provide a thick- or multi-layered mask to prevent spreading of the light on the substrate. Generally, lasers may be preferable because they can more easily provide wavelengths particularly suited for a chromophore of the photosensitive group.

While the invention is illustrated primarily herein by way of the use of a mask to illuminate the substrate, other techniques may also be used. For example, the substrate may be rotated under a modulated laser or diode light source. Such techniques are discussed in, for example, U.S. Patent No. 4,719,615, which is incorporated herein by reference.

The substrate may be irradiated either in contact with or not in contact with a solution and, preferably, is irradiated in contact with the solution. The solution may contain reagents to prevent by-products of irradiation from interfering with subsequent binding reactions. Such by-products might include, for example, carbon dioxide, nitrosocarbonyl compounds, styrene derivatives, indole derivatives, and products of their photochemical reactions. Reagents added to the solution may include, for example, acidic or basic buffers, thiols, substituted hydrazines and hydroxylamines, reducing agents (e.g., NADH or bisulfite ion) or reagents known to react with a given functional group (e.g., aryl nitroso + glyoxylic acid \rightarrow aryl formhydroxamate + CO₂).

Preferably, however, protecting groups will be selected which do not cause significant interferences with the binding reactions. Also, wash steps will be incorporated so that the by-products do not interfere with the reactions.

5 In a preferred embodiment, a surface provided with a plurality of sites occupied by photosensitive N-derivatives of biotin or biotin analogs is exposed to a desired light pattern to cause loss of some or all of the photosensitive protecting groups at predefined regions on the surface. Such irradiation
10 of the N-derivatized biotin compounds of the present invention leads to formation of surface-bound biotin or biotin analogs having a strong specific binding affinity for avidin or avidin analogs. The specific binding affinity of biotin and avidin is one of the strongest known between macromolecules ($K_a = 10^{15}$
15 M^{-1}). This binding persists when the carboxyl terminus of biotin is attached to another entity, e.g., a surface, or when avidin is attached to another molecule. Avidin possesses four subunits having specific binding affinity for biotin molecules. For example, deprotected biotin sites may be incubated with
20 avidin or an avidin conjugate of an anti-ligand, e.g., an antibody, to provide a localized concentration of the desired anti-ligand on the surface. When incubation with avidin alone is performed, it is necessary to further incubate the resulting product with a preselected species having specific binding
25 affinity for avidin, e.g., a biotinylated anti-ligand. Thus, biotinylated anti-ligands can be bound to the free sites of avidin to afford anti-ligands immobilized at predefined regions on the surface. For a general discussion of the use of the biotin-avidin interaction in molecular biology, see Bayer, et
30 al. Once localization of the anti-ligand is complete, the light pattern can be changed and the same or a different anti-ligand can be localized at other discrete sites on the surface.

35 V. Attachment of Anti-ligands

An anti-ligand is one or more molecules that recognize a particular ligand in solution. Examples of ligands that can be investigated by this invention include, but are not

restricted to agonists and antagonists for cell membrane receptors, toxins and venoms, viral epitopes, antigenic determinants, hormones, hormone receptors, steroids, peptides, enzymes, substrates, cofactors, drugs, lectins, sugars, oligonucleotides, oligosaccharides, proteins, and monoclonal and polyclonal antibodies.

Anti-ligands that mediate a biological function on binding with particular ligand(s) are of most interest. Suitable anti-ligands include relatively small, single molecules, such as cofactors, which show specific binding properties. Typically, anti-ligands will be greater than about 100 daltons in size and more typically will be greater than about 1kD in size. Other examples of anti-ligands include, but are not restricted to, the common class of receptors associated with the surface membrane of cells and include, for instance, the immunologically important receptors of B-cells, T-cells, macrophages and the like. Other examples of anti-ligands that can be investigated by this invention include but are not restricted to hormone receptors, hormones, drugs, cellular receptors, membrane transport proteins, steroids, peptides, enzymes, substrates, cofactors, vitamins, lectins, sugars, oligonucleotides, oligosaccharides, viral epitopes, antigenic determinants, glycoproteins, and immunoglobulins, e.g., monoclonal and polyclonal antibodies.

In a preferred embodiment, the anti-ligand will be a biotinylated receptor which binds specifically to avidin. Many biotinylated anti-ligands and biotinylating reagents are commercially available. (See, for example, Vector Laboratories, Inc., Catalog, Burlingame, CA) Methods for biotinylating desired anti-ligands are well-known in the art and are described, for example, at Bayer, et al.

In a preferred embodiment a plurality of anti-ligands is immobilized on a surface by first attaching photoreactive caged binding members to the surface. The caged binding members on a predefined region of the surface are exposed to light to give binding members having a high affinity for a specific binding substance. The activated binding members on the predefined region are then incubated with the specific

binding substance, the surface is washed free of unbound specific binding substance, and the surface is incubated with a desired anti-ligand or anti-ligand conjugate. The exact incubation conditions, e.g., time, temperature, pH, will depend upon the species used and will be readily apparent to one skilled in the art. After washing the surface free of unbound anti-ligand, the above steps can be repeated on a different region of the surface.

In another embodiment of the invention a plurality of anti-ligands is immobilized on a surface as described above, except the attachment of anti-ligands to specific binding substance is carried out prior to introducing the specific binding substance to the surface.

In a further embodiment the anti-ligand is a monoclonal or polyclonal antibody. In a still further preferred embodiment the anti-ligand is a biotinylated antibody or biotinylated receptor.

A most preferred embodiment of the invention is when the binding member is biotin or a biotin analog and the specific binding substance is avidin or an avidin analog.

VI. Screenings and Assays

A surface prepared according to the methods described above can be used to screen for ligands having high affinity for immobilized anti-ligands. Screening can be performed by immobilizing a plurality of anti-ligands on predefined regions of a surface by the methods described above. A solution containing a marked (labelled) ligand is introduced to the surface and incubated for a suitable period of time. The surface is then washed free of unbound ligand and the anti-ligands having high affinity for the ligand are identified by identifying those regions on the surface where markers are located. Suitable markers include, but are not limited to, radiolabels, chromophores, fluorophores, chemiluminescent moieties, and transition metals. Alternatively, the presence of ligands may be detected using a variety of other techniques, such as an assay with a labelled enzyme, antibody, and the like. Other techniques using various marker systems for

detecting bound ligand will be readily apparent to those skilled in the art.

In a preferred embodiment, a substrate prepared as discussed above can be exposed to a solution containing marked
5 ligand such as a marked antigen. The ligand can be marked in any of a variety of ways, but in one embodiment marking is effected with a radioactive label. The marked antigen binds with high affinity to an immobilized antibody previously
10 localized on the surface. After washing the surface free of unbound ligand, the surface is placed proximate to x-ray film to identify the antibodies that recognize the antigen. Alternatively, a fluorescent marker may be provided and detection may be by way of a charge-coupled device (CCD), fluorescence microscopy or laser scanning.

15 When autoradiography is the detection method used, the marker is a radioactive label, such as ^{32}P . The marker on the surface is exposed to X-ray film, which is developed and read out on a scanner. An exposure time of about 1 hour is typical in one embodiment. Fluorescence detection using a
20 fluorophore label, such as fluorescein, attached to the ligand will usually require shorter exposure times.

Quantitative assays for ligand concentrations can also be performed according to the present invention. In a direct assay method, the surface containing localized
25 anti-ligands prepared as described above, is incubated with a solution containing a marked ligand for a suitable period of time. The surface is then washed free of unbound ligand. The amount of marker present at predefined regions of the surface is then measured and can be related to the amount of ligand in
30 solution. Methods and conditions for performing such assays are well-known and are presented at, for example, L. Hood, et al., Immunology, Benjamin/Cummings (1978) and E. Harlow, et al., Antibodies. A Laboratory Manual, Cold Spring Harbor Laboratory, (1988). See, also U.S. Patent No. 4,376,110 for
35 methods of performing sandwich assays. The precise conditions for performing these steps will be apparent to one skilled in the art.

A competitive assay method can also be employed by the present invention. Such a method involves immobilizing anti-ligands on predefined regions of a surface as described above. An unmarked ligand is then bound to anti-ligands on the surface having specific binding affinity for the ligand. A solution containing marked ligand is then introduced to the surface and incubated for a suitable time. The surface is then washed free of unbound reagents and an amount of marker remaining on the surface is measured. Alternatively, marked and unmarked ligand can be exposed to the surface simultaneously. The amount of marker remaining on predefined regions of the surface can be related to the amount of unknown ligand in solution.

Use of the invention herein is illustrated primarily with reference to screenings of ligands for anti-ligands and assays for ligands. The invention will, however, find many other uses. For example, the invention may be used in information storage (e.g., on optical disks), production of molecular electronic devices, production of stationary phases in separation sciences, and in immobilization of cells, proteins, lectins, nucleic acids, polysaccharides and the like in any desired pattern on a surface via molecular recognition of a specific anti-ligand.

The invention has been described primarily with reference to the use of photoremovable protecting groups, but it will be readily recognized by those of skill in the art that other types of groups can be used and that other sources of radiation can also be used. For example, in some embodiments it may be desirable to use protecting groups sensitive to electron beam irradiation, X-ray irradiation, X-ray lithography, or combinations thereof. Alternatively, the group could be removed by exposure to an electric current.

It is to be understood that the above description is intended to be illustrative and not restrictive. Many embodiments will be apparent to those of skill in the art upon reviewing the above description. The scope of the invention should, therefore, be determined not with reference to the above description, but should instead be determined with

reference to the appended claims, along with the full scope of equivalents to which such claims are entitled.

VII. Examples

5 The following examples of preferred embodiments of the present invention are presented by way of illustration only and do not suggest that the above-described methods and compositions are in any way limited by the specific examples set forth below.

10

Synthesis of Photoreactive N-1'-Derivatives of Biotin

 Methods for the preparation of acyl imidazolinones, such as biotin derivatives, are well-known. See, for example, Kohn, et al., J. Org. Chem. (1977), 42, 941-948 and Knappe, et al., Biochem. Z. (1961), 335, 168-176. Treatment of biotin methyl ester with methyl chloroformate in refluxing chloroform (no base) for 72-80 h afforded a mixture heavily favoring the N-1'-derivative. Under similar conditions, the use of nitroveratryloxycarbonyl (NVOC) chloride (Amit, et al., J. Org. Chem. (1974), 39, 192-196) gave N-
15 1'-(nitroveratryloxycarbonyl)-biotin methyl ester (NVOC-biotin-OMe) in 47% yield after chromatography and crystallization. Likewise, N-1'-(nitroveratryloxycarbonyl)-biotin p-nitrophenyl ester
20 (NVOC-biotin-ONP) was obtained in 39% yield.

 The structural assignment is based on precedent as well as spectroscopic properties. See, for example, E. Becker, High Resolution NMR, 2nd ed., Acad. Press (1980). In particular, the ¹H NMR spectrum readily differentiates the ring
30 fusion protons bearing 1) a urea nitrogen (ca 4.2 ppm) and 2) an imide nitrogen (ca 4.8 ppm). Through the use of COSY (Derome, Modern NMR Techniques for Chemistry Research, Pergamon Press, Oxford (1987)) on N-1'-(nitroveratryloxycarbonyl)-biotin p-nitrophenyl ester, it was determined that the former ring
35 fusion proton is vicinal to a methine adjacent to sulfur, and that the latter is vicinal to a methylene adjacent to sulfur.

 Using the conditions reported by Kohn, et al. biotin methyl ester is similarly derivatized with


dimethyldimethoxycarbobenzyloxycarbonyl (DDZ) chloride or 1-pyrenylmethyloxycarbonyl (PYROC) chloride to afford the photolabile compounds N-

1'-(dimethyldimethoxycarbobenzyloxycarbonyl) biotin methyl ester (DDZ-biotin-OMe) and N-1'-(1-pyrenylmethyloxycarbonyl) biotin methyl ester (PYROL-biotin-OMe), respectively. The methods of Knappe, et al., and Kohn, et al., can be employed to prepare analogous compounds and are incorporated by reference herein.

10

Example A: Preparation of NVOC-biotin-OMe
(N-1'-(6-nitroveratryloxycarbonyl)-biotin methyl ester)

15



20

2.00 g (8.19 mmol) of D-biotin was added to a methanolic HCl solution prepared from 2.5 ml of acetyl chloride in 40 ml of anhydrous methanol. After stirring for 15 hours, the solvent was removed under reduced pressure to afford 2.11 g of the product biotin-OMe (biotin methyl ester) as a white solid, MP 116-118° C (100% yield).

NVOC-biotin-OMe was prepared from biotin via the intermediate, biotin methyl ester (biotin-OMe) by either of two methods.


1. A solution of 1.00 g (3.87 mmol) of biotin-OMe and 1.60 g (5.81 mmol) of 6-nitroveratryloxycarbonyl chloride in 10 ml of chloroform was heated to reflux for 50 hours. The product was purified via flash-column chromatography on silica gel (3% methanol, 3% acetone, 94% chloroform as eluent) to afford 0.90 g of NVOC-biotin-OMe as a yellow solid (47% yield, 84% yield based on unreacted starting biotin-OMe), MP

199-203° C, and 0.44 g of recovered biotin-OMe (44% recovery). The product, NVOC-biotin-OMe, was recrystallized from methylene chloride/ether.

2. A solution of 3.4 g (12 mmol) of
5 6-nitroveratryloxycarbonyl chloride in 40 ml of methylene chloride was added to a solution of 1.1 ml (14 mmol) of pyridine and 1.3 g (13 mmol) of phenol in 15 ml of methylene chloride cooled to 0° C. The reaction mixture was allowed to warm to room temperature and was stirred for 19 hours. The
10 solution was partitioned between methylene chloride and 1 N HCl, the organic phase was separated and dried with magnesium sulfate, and the solvent was removed under reduced pressure to give 4.1 g of a brown oil. Purification via flash-column chromatography on silica gel (90% methylene chloride/10% hexane
15 as the eluent) afforded 2.0 g of the product, 6-nitroveratryl phenyl carbonate, as a colorless oil.


A mixture of 399 mg (1.20 mmol) of 6-nitroveratryl phenyl carbonate in 5 ml of chloroform and 202 mg (0.782 mmol) of biotin-OMe were heated to reflux. After 50 hours TLC showed
20 that no reaction occurred, therefore 35 mg of 60% sodium hydride (0.88 mmol) was added in two equal portions over 15 minutes. After an additional 16 hours at reflux, the reaction was quenched with 3 drops of glacial acetic acid. The product was purified via flash-column chromatography on silica gel (3%
25 methanol, 3% acetone, 94% chloroform as the eluent) to afford 264 mg of the product, NVOC-biotin-OMe, as an off-white solid (68% yield, 75% yield based on recovered unreacted biotin-OMe) and 20 mg of recovered biotin-OMe (10% yield).

Example B: Preparation of NVOC-biotin-ONP
(N-1'-(6-nitroveratryloxycarbonyl)-biotin
para-nitrophenyl ester)



A mixture of 0.340 g (0.930 mmol) of biotin
p-nitrophenyl ester (purchased from Sigma Chemical Co., St.
Louis) and 0.450 g (1.63 mmol) of 6-nitroveratryloxycarbonyl
chloride in 4 ml of chloroform was heated to reflux for 65
hours. The mixture was purified via flash-column
chromatography on silica gel (3% methanol, 3% acetone, 94%
chloroform as the eluent) to produce 0.231 g of NVOC-biotin-ONP
as a beige solid (39% yield, 93% yield based on unreacted
biotin-ONP), MP 203-205° C, and 0.21 g of recovered unreacted
biotin-ONP (58% yield). The product was further purified via
recrystallization from chloroform/hexane.


Example C: Preparation of NPOC-biotin-ONP (N-1'-(6-
nitropiperonyloxycarbonyl)-biotin para-
nitrophenyl ester)



A mixture of 0.365 g (1.00 mmol) of biotin p-
nitrophenyl ester and 0.410 g (1.58 mmol) of 6-

nitropiperonyloxycarbonyl chloride in 5 ml of chloroform was heated at reflux for 62 hours. The product was purified via flash-column chromatography on silica gel (3% methanol, 3% acetone, 94% chloroform as the eluent) to produce 0.231 g of product as a beige solid (39% yield, 93% based on unrecovered starting material), MP 203-205° C, and 0.21 g of recovered biotin-ONP (58% yield). As described, the product was further purified via recrystallization from chloroform/hexane.

10 Example D: Preparation of NVOC-DT-biotin-OMe
(N-1'-(6-nitroveratryloxycarbonyl)-dethiobiotin
methyl ester)




20

A slurry of 50 mg (0.19 mmol) of biotin-OMe and about 1 g of Raney Nickel active catalyst (50% solution in H₂O from Aldrich Chemical Co.) in 4 ml of methanol was stirred at room temperature for one hour. The reaction mixture was diluted with chloroform, filtered and the recovered catalyst was washed with methanol. The combined wash and filtrate were partitioned between chloroform and saturated sodium chloride acidified to pH 2 with 1 N HCl. The combined organic phases were dried over magnesium sulfate, and the solvent removed under reduced pressure to give 34 mg of pure DT-biotin-OMe as a white solid, MP 68-72° C (77% yield).

A mixture of 34 mg (0.15 mmol) of DT-biotin methyl ester and 74 mg (0.27 mmol) of 6-nitroveratryloxycarbonyl chloride in 3 ml of chloroform was heated to reflux for 15 hours. The products are purified via flash-column chromatography on silica gel (3% methanol, 97% chloroform as

the eluent) to afford 45 mg of a 1:3 mixture of products as a yellow solid (65% yield), MP 152-156° C.

Example E: Preparation of NVOC-biotin-OH (N-1'-(6-nitroveratryloxycarbonyl)-biotin)



A solution of 262 mg (0.527 mmol) of NVOC-biotin-OME, prepared in by either method of Example A, in 15 ml of tetrahydrofuran and 2 ml of dimethylformamide was treated with 10 ml of 1 N HCl. The reaction mixture was heated to reflux for 49 hours, cooled to room temperature, and the solvent removed under reduced pressure. The crude product was purified via flash-column chromatography on silica gel (10% methanol, 90% chloroform as the eluent) to afford 178 mg of the pure product, NVOC-biotin-OH, as a white solid (70% yield), MP 219-223° C.

Example F: Chromatographic Evidence for Photoremoval of the NVOC group from NVOC-biotin-ONP

A 100 mM solution of NVOC-biotin-ONP in acetonitrile, or biotin in water, was placed in a quartz cuvette with a 2.0 mm pathlength. The cuvette was irradiated for two minutes at a power of 1 watt/cm² with a 500 W Hg(Xe) arc lamp (Oriel #66142) having a 305 nm long pass filter (Oriel #51450). Illuminated and non-illuminated samples were then subjected to reverse-phase HPLC.

Shown in FIG. 1 are chromatographs of illuminated biotin and illuminated NVOC-biotin-ONP. The results support the following: 1) biotin was unaffected by the illumination;

dimethylsulfoxide to a solution of 43 mg (0.090 mmol) of NVOC-biotin-OH, 13 mg (0.096 mmol) of HOBt, and 0.035 ml of diisopropylethyl amine in 0.090 ml of dimethylsulfoxide. After five minutes, the resulting solid was diluted to 1.80 ml with dimethylsulfoxide to give a 0.05 M solution of the activated ester. Approximately 0.5 ml of the activated ester solution was applied to the surface of the derivatized glass surface. After being exposed to the activated ester for two hours, the slides were washed sequentially with dimethylformamide, ethanol, and methylene chloride to yield a NVOC-biotin-caproic-propyl derivatized surface.

Method B (via the ONP ester)

Approximately 0.5 ml of an 0.10 M solution of NVOC-biotin-ONP in dimethylformamide was applied to the surface of the derivatized glass slide. After 2-24 hours, the slides were washed sequentially with dimethylformamide, ethanol, and methylene chloride to yield a NVOC-biotin-caproic-propyl derivatized surface.

Example J: Photodeprotection of an NVOC-Biotin-Caproic-Propyl-derivatized slide, and subsequent labeling with a Fluorescein-Streptavidin conjugate

A glass microscope slide to which NVOC-biotin has been covalently attached via a caproic-propyl spacer, as described in the Example I, was mounted on a custom flow cell and illuminated through a 500 μm x 500 μm checkerboard-pattern mask (Photo Sciences Inc., Torrance, CA) using broad-band UV/blue light. The light source was a 500 W Mercury arc lamp (Oriel Model 87330) equipped with a 350 nm - 450 nm dichroic reflector and produced actinic light having an intensity of 12 mW/cm² as measured through a 360 nm bandpass filter. The derivatized surface was photolyzed in flowing dioxane for 15 minutes, removed from the flow cell, and sequentially rinsed in deionized water, ethanol, and methylene chloride.

After incubation for one hour in a solution containing filtered PBS, 1% BSA, 0.05% Tween 20, pH 7.4, the activated surface was treated with a solution containing a Fluorescein-derivative of Streptavidin (Molecular Probes; 10 μ g/ml in PBS/BSA/Tween 20) for one hour at room temperature. The slide was vortexed twice in PBS, 0.05% Tween 20, pH 7.4 for 10 minutes, rinsed with deionized water, and allowed to dry. The slide was examined with a scanning fluorescence microscope (Zeiss Axioskop equipped with a Newport Model PM500-C motion controller, a Spectra-Physics Model 2020 argon-ion laser producing a 488 nm excitation light; and a 520 nm long-pass emission filter) interfaced with a photon-counting device (Hamamatsu Model 9403-02 photomultiplier; Stanford Research Systems Model SR445 amplifier and Model SR430 multichannel scaler, IBM compatible PC) to generate a two-dimensional image consisting of fluorescence intensity data as a function of x,y position. An example of this technique is described in U.S. Patent Applications Serial Numbers 07/362,901 and 07/492,462, filed June 7, 1989 and March 7, 1990, respectively, and incorporated herein by reference. FIG. 4 shows an example of the images obtained.

The light squares indicate regions of high fluorescence intensity resulting from localization of the fluorescein label attached to the anti-ligand. This experiment demonstrates enhanced binding of streptavidin upon photodeprotection of NVOC-biotin coupled to a surface with a caproic-propyl spacer, and spatially-addressable immobilization of an anti-ligand, such as streptavidin, by non-covalent means.

Example K: Photodeprotection of an NVOC-Biotin-Caproic-Propyl-derivatized slide, treatment with Streptavidin, and labeling with a Fluorescein-Biotin conjugate

The microscope slide of Example I having NVOC-biotin covalently attached via a caproic-propyl spacer was illuminated and processed as in Example J, except that after preincubation with the PBS/BSA/Tween 20 solution, the surface was treated with 10 μ g/ml solution of Streptavidin (in PBS/BSA/Tween 20)

for 40 minutes at room temperature, followed by incubation with a 1 μ M solution of Fluorescein-Biotin (5-(N-((5-(N-(6-(biotinoyl)amino)hexanoyl)amino)pentyl)thioureidyl)fluorescein, in PBS, pH 7.4) for 20 minutes. The resulting slide was then washed, dried, and examined using a scanning fluorescence microscope as described above. FIG. 5 shows an example of the images obtained.

The light squares indicate regions of high fluorescence intensity resulting from localization of the fluorescein label attached to ligand, biotin. This experiment demonstrates the binding of a ligand (a Fluorescein-Biotin complex) to streptavidin immobilized in a spatially-addressable manner.

15

Example L: Photodeprotection of an NVOC-Biotin-Polyether-derivatized slide and labeling with a Fluorescein-Streptavidin conjugate

A microscope slide to which 3-aminopropyltriethoxy silane had been attached was treated with the BOP-activated ester of 18-amino-6-aza-10,15-dioxa-5-ketooctadecanoic acid using a procedure similar to Example I. The resulting slide was then derivatized with NVOC-Biotin-ONP, and was illuminated, processed, and examined as described in Example J. FIG. 6 shows an example of the images obtained.

25

This experiment demonstrates spatially-localized streptavidin binding upon photodeprotection of NVOC-Biotin coupled to a surface using an alternative linker, a polyether-glutaric-propyl moiety.

30

Example M: Photodeprotection of an NPOC-Biotin-Caproic-Propyl-derivatized slide and labeling with a Fluorescein-Streptavidin conjugate

The microscope slide to which NPOC-biotin-ONP had been covalently attached via a caproic-propyl spacer was illuminated, processed, and examined as described in Example J. FIG. 7 shows an example of the images obtained.

35

This experiment demonstrates spatially-localized streptavidin binding upon photodeprotection of caged-biotin using a different protecting group, NPOC.

5 Example N: Photodeprotection of an
 NVOC-Biotin-Caproic-Propyl-derivatized slide in
 aqueous buffer and labeling with a
 Bodipy-streptavidin conjugate

10 The microscope slide of Example I having NVOC-Biotin
covalently attached via a caproic-propyl spacer was mounted on
a custom flow cell and illuminated using the apparatus
described in Example J. The derivatized surface was photolyzed
in PBS, 1% BSA, 0.1% Tween 20 for 30 minutes at 12 mW/cm²,
removed from the flow cell, and processed as described in
15 Example K, except that Bodipy-streptavidin (Molecular Probes,
50 µg/ml) was used instead of the fluorescein conjugate. FIG. 8
shows an example of the images obtained.

 This experiment demonstrates spatially-localized
streptavidin binding upon photodeprotection of NVOC-biotin in a
20 different solvent, in this case an aqueous buffer.

 Example O: Evaluation of crosslinking groups of different
 lengths

 A microscope slide derivatized with
25 N-Boc-3-aminopropyltriethoxy silane, as in Example I, was
functionalized with three different crosslinkers in different
locations by selectively coupling 1, 2 or 3
N-Boc-6-aminocaproic acid linkers to the surface via their BOP-
activated esters. This generated four distinct and well
30 defined regions on the surface of the slide having zero, one,
two and three 6-aminocaproic acid linkers (4, 11, 18 and 25
atom spacers, respectively) between the surface and the
terminal amino functionality used to bind to the carboxyl group
of the derivatized biotin. A 0.1 M solution of biotin
35 p-nitrophenyl ester in dimethylformamide was subsequently
coupled to the slide, and the relative binding affinity of
streptavidin to the surface-bound biotin was measured by
incubating the slide with fluoresceinated streptavidin and

measuring the fluorescence intensity as in Example J. The measured relative fluorescence was 38, 68, 85 and 100% (normalized to the fluorescence of area having three caproic linkers), respectively, for zero, one two and three caproic linkers, showing that a higher density of streptavidin was bound to an area of the slide derivatized with biotin spaced relatively far from the surface of the slide. FIG. 9 shows the fluorescence of fluoresceinated streptavidin bound to the glass slide derivatized in this experiment.

10

Example P: Photodeprotection of an NVOC-Biotin-Caproic-Propyl-derivatized slide and subsequent immobilization of antibodies on the derivatized surface.

15

A microscope slide to which NVOC-Biotin had been covalently attached via a caproic-propyl spacer was mounted on a custom flow cell and illuminated as described in Example J, except that a hand-cut mask consisting of a horizontal stripe (approximately 2 mm wide) was used. After removal from the flow cell and rinsing, the slide was incubated for 30 minutes with PBS, 1% BSA, 0.05% Tween 20, pH 7.4, followed by 30 minutes treatment with Streptavidin (10 µg/ml in PBS/BSA/Tween 20), rinsing with PBS, 0.05% Tween 20, 60 minutes incubation with biotinylated Rabbit IgG (Vector Laboratories; 50 µg/ml in PBS/BSA/Tween 20), and rinsing with PBS/Tween 20. The surface was then "capped" to prevent subsequent streptavidin binding to the multiple biotin moieties on the biotinylated IgG. That is, streptavidin was used to bind free biotin on IgG. This was accomplished by re-treatment with Streptavidin solution, rinsing with PBS/Tween 20, followed by incubation with free biotin (Molecular Probes; 1 mM in PBS/Tween 20; 10 minutes), and a final PBS/Tween 20 rinse. The slide was then remounted on the flow cell, photolyzed for 30 minutes in PBS/Tween 20 using a hand-cut mask consisting of vertical stripes (approximately 2 mm wide), and processed as described above, except that biotinylated Mouse IgG (Vector Laboratories; 50 µg/ml in PBS/BSA/Tween 20; 30 minute incubation) was used, and the "capping" steps were not repeated. The slide was then

30
35

rinsed with deionized water, allowed to dry, and beads of silicone gasket compound (Permatex Ultra Blue) were used to partition the slide into three areas. After pre-incubation for 30 minutes with PBS/BSA/Tween 20, the first area was treated with Fluorescein-labeled anti-Rabbit IgG (Vector Laboratories; made in goat; 100 μ g/ml in PBS/BSA/Tween 20). The second area was treated with Fluorescein-labeled anti-Mouse IgG (Vector Laboratories; made in horse; 100 μ g/ml in PBS/BSA/Tween 20). The third area was incubated with an equimolar mixture of the two secondary antibodies. The slide was then vortexed twice in PBS/Tween 20 for two minutes, rinsed briefly with deionized water, and allowed to dry. The different regions of the slide were examined using the scanning fluorescence microscope described in Example J.

FIG. 10 shows examples of the images obtained from the glass slide derivatized in this experiment. Figure 10a shows the region of the glass slide treated with fluorescein-anti-Rabbit IgG. As expected, the horizontal stripe, which corresponds to the area where biotinylated Rabbit IgG bound, is intensely fluorescent indicating a high density of bound fluorescein-anti-Rabbit IgG. The vertical stripes in this region are faintly visible, which may be due to the slight cross reactivity of the secondary antibodies. FIG. 10b shows the region of the glass treated with Fluorescein-anti-Mouse IgG. Here, the vertical stripes, where Mouse IgG is bound, are fluorescent while the horizontal areas do not fluoresce appreciably. Finally, FIG. 10c shows the region treated with both secondary antibodies (fluorescein-anti-Rabbit and anti-Mouse). In this case both the vertical and horizontal stripes fluoresce, indicating a high surface density fluorescein and, therefore, of the secondary antibodies. This experiment demonstrates the spatially addressable immobilization of two different antibodies on the same surface.

WHAT IS CLAIMED IS:

1. A method for forming predefined regions on a
5 surface of a solid support, the predefined regions capable of
selectively immobilizing an anti-ligand, the method comprising:
 - a) attaching to the surface a caged binding member
having a low affinity for the anti-ligand and which member is
convertible by irradiation to a binding member capable of
10 immobilizing the anti-ligand by an interaction having an
affinity constant of 10^7 or stronger; and
 - b) selectively irradiating the predefined regions of
the surface to convert the caged binding member in the
predefined regions to the binding member.
- 15 2. A method as in claim 1 wherein the anti-ligand is
attached to the binding member by a specific binding substance.
3. A method as in claim 2 wherein the specific
20 binding substance is avidin or streptavidin.
4. A method as in claim 3 wherein the specific
binding substance is egg avidin.
- 25 5. A method as in claim 1 wherein the binding member
is attached to the surface by a crosslinking group.
6. A method as in claim 1 wherein the binding member
is biotin.
- 30 7. A method as in claim 1 wherein the anti-ligand is
an immunoglobulin.
8. A method as in claim 1 wherein the immunoglobulin
35 is a monoclonal antibody.

9. A method as in claim 2 wherein the binding member binds to a specific binding substance with an affinity constant (K_a) of about 10^{15} M^{-1} .

10. A method as in claim 1 wherein the surface is silica or glass.

11. A method for attaching an anti-ligand to a surface comprising the steps of:

a) linking to the surface a caged biotin molecule having a relatively low affinity for avidin;

b) irradiating caged biotin molecules on predefined regions of the surface to form biotin analogs having high binding affinities for avidin;


c) reacting the biotin analogs with avidin; and

d) incubating biotinylated anti-ligands with the predefined regions on the surface.

12. A method as in claim 11 wherein the anti-ligand is a monoclonal antibody.


13. A method as in claim 11 wherein the surface is glass or silica.

14. A method as in claim 11 wherein the caged biotin molecule has the formula:



wherein X is selected from the group consisting of oxycarbonyls of lower alkyl, aryl, and benzyl groups, provided that X is not methyl oxycarbonyl; R is selected from the group consisting of hydrogen, lower alkyl, aryl, benzyl, and N-succinimidyl; and acid addition salts of the molecule.

15. A method as in claim 14 wherein the biotin analog has the formula:



10 wherein W is a surface or a crosslinking group attached to a surface.

16. A method as in claim 11 wherein the biotin analog and avidin have an affinity constant (K_a) of about
15 10^{15} M^{-1} .

17. A method for localizing an anti-ligand on a surface comprising the steps of:

- 20 a) linking to the surface a photoactivatable biotin molecule having low affinity for avidin;
b) exposing the photoactivatable biotin molecule to light to form a biotin analog having a high binding affinity for avidin; and
c) incubating an anti-ligand attached to avidin with
25 the biotin analog.

18. A method as in claim 17 wherein the anti-ligand is attached to avidin by a bifunctional crosslinking group.


30 19. A method as in claim 17 wherein the anti-ligand is connected to avidin by a biotin group.

20. A method as in claim 17 wherein the anti-ligand is a monoclonal antibody.

35

21. A method as in claim 17 wherein the surface is glass or silica.

22. A method as in claim 17 wherein the photoactivatable biotin molecule has the formula:



10 wherein X is selected from the group consisting of oxycarbonyls of lower alkyl, aryl, and benzyl groups, provided that X is not methyl oxycarbonyl; R is selected from the group consisting of hydrogen, lower alkyl, aryl, benzyl, and N-succinimidyl; and acid addition salts of the molecule.


15

23. A method as in claim 17 wherein the anti-ligand is attached to egg avidin.

24. A method as in claim 17 wherein the biotin analog and avidin have an affinity constant (K_a) of about 10^{15} M^{-1} .

20

25. A compound having the formula:



25

30

wherein X and Z are selected from the group consisting of hydrogen and oxycarbonyls of lower alkyl, aryl, and benzyl groups, provided that when X is hydrogen, Z is not hydrogen or methyloxycarbonyl, and provided that when Z is hydrogen, X is not hydrogen or methyloxycarbonyl; R is selected from the group consisting of hydrogen, lower alkyl, aryl, lower alkyl formate, aryl formate, formamide, N-alkylformamide, N-succinimidyl, hydroxyl, alkoxyl, thiol, thioether, disulfide, hydrazide and


35

amine groups, provided that when X or Z is methyloxycarbonyl, R is not methyl formate; U is O, S, or NH; Y is selected from the group consisting of sulfur, oxygen, methylene, carbonyl, sulfinyl and sulfonyl groups or Y represents two hydrogen atoms attached to the respective carbons; n = 0-7; and acid addition salts of the compound.

26. A compound as in claim 25 wherein U is O, Y is sulfur, and n = 4.

27. A compound as in claim 25 wherein one of X and Z is a nitroveratryloxycarbonyl group.

28. A compound as in claim 25 wherein one of X and Z has the formula:



wherein R₁ and R₂ are selected from the group consisting of hydrogen, lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxy, thiol, thioether, amino, nitro, carboxyl, formate, formamido and phosphido groups.


29. A compound as in claim 25 wherein X is 6-nitroveratryloxycarbonyl; Z is hydrogen; and R is methyl formate.

30. A compound as in claim 25 wherein X is 6-nitroveratryloxycarbonyl; Z is hydrogen; and R is p-nitrophenyl formate.

31. A compound as in claim 25 wherein X is 6-nitropiperonyloxycarbonyl, Z is hydrogen; and R is p-nitrophenyl formate.

32. A compound as in claim 25 wherein one of X and Z is a ring-disubstituted benzyloxycarbonyl group.

33. A compound as in claim 32 wherein the ring-disubstituted benzyloxycarbonyl group has the formula:




10

wherein R_1 , R_2 , R_3 , and R_4 are selected from the group consisting of hydrogen, lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxyl, thiol, thioether, amino, nitro, carboxyl, formate, formamido and phosphido groups.

34. A compound as in claim 33 wherein R_1 and R_2 are methoxy.

35. A compound as in claim 33 wherein R_3 and R_4 are methyl.

36. A compound having the formula:



30


wherein X is selected from the group consisting of hydrogen, lower alkyl, aryl, and benzyl groups; U and W are selected from the group consisting of hydrogen, lower alkyl, aryl, and benzyl groups, provided that only one of U and W is present; R is selected from the group consisting of hydrogen, lower alkyl,

35

aryl, carboxylate, alkyl formate, aryl formate, formamide, N-alkyl formamide, N-succinimidyl, hydroxyl, alkoxyl, thiol, thioether, disulfide, hydrazide, and amine groups; Y is selected from the group consisting of sulfur, oxygen, methylene, carbonyl, sulfinyl and sulfonyl groups, or Y represents two hydrogens attached to the respective carbons; n = 0-7; and acid addition salts of the compound.

37. A compound as in claim 36 wherein Y is sulfur and n = 4.

38. A compound as in claim 36 wherein the ring-disubstituted benzyl group has the formula:




wherein R_1 and R_2 are selected from the group consisting of hydrogen, lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxyl, thiol, thioether, amino, nitro, carboxyl, formate, formamido and phosphido groups.

39. A compound as in claim 38 wherein R_1 and R_2 are methoxy.

40. A compound as in claim 38 wherein R_3 and R_4 are methyl.

41. A compound as in claim 36 wherein X is a nitroveratryl group.

42. A compound as in claim 36, wherein X has the formula:



wherein R₁ and R₂ are selected from the group consisting of hydrogen, lower alkyl, aryl, benzyl, halogen, hydroxyl, alkoxyl, thiol, thioether, amino, nitro, carboxyl, formate, formamido and phosphido groups.

43. A compound as in claim 36 wherein X is 6-nitroveratryl; U or W is hydrogen; and R is methyl formate.

44. A compound as in claim 36 wherein X is 6-nitroveratryloxycarbonyl hydrogen; and R is p-nitrophenyl formate.

45. A compound as in claim 36 wherein X is 6-nitropiperonyloxycarbonyl and U or W is hydrogen.

46. A compound as in claim 36 wherein X is a ring-disubstituted benzyl group, and R is methyl formate or p-nitrophenyl formate.

47. A composition comprising a compound of claim 25 covalently attached to a glass or silica surface.

48. A method for localizing a plurality of anti-ligands having different ligand binding specificities on a surface comprising the steps of:

- a) attaching caged binding members to the surface;
- b) exposing the caged binding members on a first predefined region of the surface to an energy source, thereby

forming binding members on the first predefined region having a strong affinity for a specific binding substance;

c) reacting the binding members on the first predefined region with the specific binding substance;

5 d) washing the surface to remove unbound specific binding substance;

e) exposing the surface to an anti-ligand;

f) washing the surface free of unbound anti-ligand;

and

10 g) repeating steps (b-f) on a different region of the surface with a different anti-ligand.

49. A method for localizing a plurality of anti-ligands having different ligand binding specificities on a surface comprising the steps of:

15 a) attaching caged binding members to the surface;

b) exposing the caged binding members on a first predefined region of the surface to an energy source, thereby producing binding members on the first predefined region having

20 a strong affinity for a specific binding substance;

c) reacting the binding members on the first predefined region with an anti-ligand attached to a specific binding substance;

d) washing the surface to remove unbound anti-ligand;

25 and

e) repeating steps (b)-(d) on a different region of the surface with a different anti-ligand.

50. A method according to claim 48 wherein the energy source is ultraviolet light.

51. A method for screening a plurality of anti-ligands localized on predefined regions of a surface for affinity for ligands in solution comprising:

35 a) exposing the surface to a solution containing at least one marked ligand;

b) washing the surface free of unbound ligand; and





c) identifying ligands binding to anti-ligands by determining regions on the surface where markers are located.

52. A method for performing a direct assay of a
5 marked ligand in solution by measuring the ligand's affinity for anti-ligands immobilized on predefined regions of a surface comprising the steps of:
a) exposing the marked ligand to the surface; and
b) measuring an amount of the ligand attached to the
10 predefined regions of the surface.

53. A method for performing a competitive assay of a ligand in solution by measuring the ligand's affinity for anti-ligands immobilized on predefined regions of a surface
15 comprising the steps of:
a) incubating a solution containing one or more marked ligands and unknown ligands with the surface;
b) washing the surface free of unbound ligands; and
c) measuring an amount of marked ligand remaining on
20 the predefined regions of the surface.

54. A method for performing a competitive assay of a ligand in solution by measuring the ligand's affinity for anti-ligands immobilized on predefined regions of a surface
25 comprising the steps of:
a) binding unknown ligands to anti-ligands on the surface;
b) incubating a solution containing one or more marked ligands with the surface;
30 c) washing the surface free of unbound ligands; and
d) measuring an amount of marked ligand remaining on the predefined regions of the surface.

1 / 6



2 / 6




FIG. 2.

3 / 6




FIG. 3.

4 / 6




FIG._4.



FIG._5.




FIG._6.




FIG._7.

5 / 6





FIG. 8.



REPLICATE 1

REPLICATE 2


REPLICATE 3

REPLICATE 4

3 2 1 0

CAPROIC ACID LINKERS

FIG. 9.



INTERNATIONAL SEARCH REPORT

International Application No. PCT/US90/06607

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC(5): A01N 1/02; C12 Q 1/00; G01N 33/566, 33/543 ;007D 471/02,235/00,473/00,235/30 U.S. CL.: 427/2;430/270,343,371,541;435/7.5;436/501,518,527;548/113,302,303,323		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
U.S. CL.	427/2; 430/270, 343, 371, 541; 435/7.5; 436/501, 518, 527; 548/113, 302, 303, 323	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ⁹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	U.S. A. 4.557.998 (Washburn et al.) 10 December 1985. See column 2-column 3 and column 12. lines 49-59.	1.2.5.13. 48-50
Y	U.S. A. 4.722.906 (Guire) 02 February 1988. See column 2-column 4.	1-54
Y	U.S. A. 4.567.902 (Saenger et al) 18 March 1986. See column 2.	25-47
X	U.S. A. 4.709.044 (Sklavounos) 24 November 1987. See column 2-column 3.	14-35
X	U.S. A. 4.016.043 (Schuurs et al.) 05 April 1977. See column 2-column 3.	51-54
X	U.S. A. 4.371.515 (Chu) 01 February 1983. See column 1-column 2.	51-54
X.P	U.S. A. 4.895.809 (Schlabach et al.) 23 January 1990. see entire document.	51-54
<p>* Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Δ" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
31 January 1991	11 MAR 1991	
International Searching Authority	Signature of Authorized Officer	
ISA/US	Jacintha Stall	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category *	Citation of Document, with indication, where appropriate, of the relevant passages.	Relevant to Claim No
Y	Journal Of Organic Chemistry, Volume 39. No. 2. issued 1974. Amit et al.. "Photosensitive Protecting Groups of Amino Sugars and Their Use in Glycoside Synthesis. 2-Vitrobenzyloxy carbonylamino and 6-Vitroveratryloxy carbonylamino Derivatives", pages 192-196. See entire document.	25-47
Y	Journal Of The American Chemical Society. Volume 92. issued 1970. Patchornik et al.. "Photosensitive Protecting Groups". pages 6333-6334. See entire document.	25-47
Y	Annual Review Of Biophysics and Biophysical Chemistry, Volume 18. issued 1989. McCray et al.. "Properties and Uses of Photoreactive Caged Compounds". pages 239-270. See entire document.	1-54
A	Guire, "Methods In Enzymology". published 1976 by Academic Press (N.Y.). See pages 280-288.	1-54
A	Journal Of Organic Chemistry, Volume 42. No. 6. issued 1977. Kohn et al.. "Syntheses and Spectral Properties of Substituted Imidazolidones and Imidazolines". pages 941-948.	25-48
A	Biochemical And Biophysical Research Communication, Volume 136, No. 1. issued 14 April 1986. Roffman et al.. "Selective Labeling of Functional Groups on Membrane Proteins or Glycoproteins Using Reactive Biotin Derivatives and ¹²⁵ I-Streptavidin". pages 80-85.	1-54
A	Analytical Biochemistry, Volume 163. issued 1987, Lacey et al., "Photobiotin as a Sensitive Probe for Protein Labeling", pages 151-158.	1-50

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category *	Citation of Document, with indication, where appropriate, of the relevant passages.	Relevant to Claim No
A	Journal Of The American Chemical Society. Volume 100. No. 11. issued 24 May 1978. Hofmann et al.. "Avidin-Biotin Affinity Columns. General Methods for Attaching Biotin to Peptides and Proteins". pages 3585-3590.	1-54
A	Journal Of Polymer Science. Volume 22. issued 1984, Ichimura. "A convenient Photochemical Method to-Immobilize Enzymes; pages 2817-2828.	1-50
A	Biochemistry. Volume 21. issued 1982. Hofmann. et al.. "Avidin Binding of Carboxyl-substituted Biotin and Analogues". pages 978-984.	1-54
A	Nucleic Acids Research. Volume 13. No. 3. issued 1985. Forster. et al.. "Non-radioactive Hybridization Probes Prepared by the Chemical Labelling of DNA and RNA with a Novel Reagent, Photobiotin." pages 745-761.	1-54
A	US. A. 4,775,745 (Ford et al.) 04 October 1988.	25-47
A.P	US. A. 4,898,951 (Symons) 06 February 1990.	25-47
A	US. A. 4,656,252 (Giese) 07 April 1987.	1-50

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE¹

This International search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers _____, because they relate to subject matter¹² not required to be searched by this Authority, namely:

2. ☐ Claim numbers _____, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out¹², specifically:

3. ☐ Claim numbers _____, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 5.4(a).

VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING³

This International Searching Authority found multiple inventions in this International application as follows:

I. A method for forming a predefined region on a surface of a solid support, a compound used in forming said predefined region, and a method of using said solid support classified in class 436, subclass 527 (claims 1-35 and 48-54).

II. A second compound, classified in class 548, subclass 323 (claims 36-46).

III. A third compound, classified in class 427, subclass 2. (claim 47)
☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application. *Telephone practice*

2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
☐ No protest accompanied the payment of additional search fees.